




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 7 :</b> <b>C12N 15/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/20571</b> <b>(43) International Publication Date:</b> 13 April 2000 (13.04.00)
<b>(21) International Application Number:</b> PCT/US99/23081 <b>(22) International Filing Date:</b> 5 October 1999 (05.10.99)  <b>(30) Priority Data:</b> 60/103,294 6 October 1998 (06.10.98) US 60/107,201 5 November 1998 (05.11.98) US  <b>(71) Applicants (for all designated States except US):</b> PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50319 (US). AGRICULTURAL GENETIC ENGINEERING RESEARCH INSTITUTE (AGERI) [EG/EG]; Agricultural Research Center, 9 Gamaa Street, Giza-12619 (EG).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BARBOUR, Eric [CA/US]; 6700 N.W. Timberline Drive, Des Moines, IA 50313 (US). MEYER, Terry, EuClaire [US/US]; 4338-101st Street, Urbandale, IA 50322 (US). SAAD, Mohammed, Eid [EG/EG]; Agricultural Genetic Engineering Research Center (AGERI), Agricultural Research Center, 9 Gamaa Street, Giza-12619 (EG).  <b>(74) Agents:</b> SPRUILL, W., Murray et al.; Alston & Bird LLP, P.O. Drawer 34009, Charlotte, NC 28234-4009 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> NOVEL MAIZE PROMOTERS    <b>(57) Abstract</b>  The present invention provides compositions and methods for regulating expression of heterologous nucleotide sequences in a plant. Compositions are novel nucleotide sequences for tissue-independent or constitutive promoters for genes encoding actin-2, enolase, Gos-2 and L41. A method for expressing a heterologous nucleotide sequence in a plant using the promoter sequences disclosed herein is provided. The method comprises transforming a plant cell with a nucleotide sequence operably linked to one of the promoters of the present invention and regenerating a stably transformed plant from the transport plant cell.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## NOVEL MAIZE PROMOTERS

## CROSS-REFERENCE TO RELATED APPLICATIONS

- 5 This application claims the benefit of U.S. Provisional Application No. 60/103,294, filed October 6, 1998 and U.S. Provisional Application No. 60/107,201 filed November 5, 1998.

## FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

- 10 This invention was made with Government support under a grant from the United States Agency for International Development (USAID), Grant Number 263-0240-G-00-6014-00. The Government has certain rights to this invention.

## FIELD OF THE INVENTION

- 15 The present invention relates to the field of plant molecular biology, more particularly to regulation of gene expression in plants.

## BACKGROUND OF THE INVENTION

- Expression of heterologous DNA sequences in a plant host is dependent  
20 upon the presence of an operably linked promoter that is functional within the plant host. Choice of the promoter sequence will determine when and where within the organism the heterologous DNA sequence is expressed. Where expression in specific tissues or organs is desired, tissue-preferred promoters may be used. In contrast, where gene expression in response to a stimulus is desired, inducible  
25 promoters are the regulatory element of choice. Where continuous expression is desired throughout the cells of a plant, constitutive promoters are utilized. Additional regulatory sequences upstream and/or downstream from the core promoter sequence may be included in expression constructs of transformation vectors to bring about varying levels of expression of heterologous nucleotide  
30 sequences in a transgenic plant.

Frequently it is desirable to have constitutive or inducible expression of a DNA sequence throughout the cells of an organism in a tissue-independent manner. For example, increased resistance of a plant to infection by soil- and air-

borne pathogens might be accomplished by genetic manipulation of the plant's genome to comprise a constitutive promoter operably linked to a heterologous pathogen-resistance gene such that pathogen-resistance proteins are continuously expressed throughout the plant's tissues.

5           Alternatively, it might be desirable to inhibit expression of a native DNA sequence within a plant's tissues to achieve a desired phenotype. In this case, such inhibition might be accomplished with transformation of the plant to comprise a constitutive, tissue-independent promoter operably linked to an antisense nucleotide sequence, such that constitutive expression of the antisense sequence  
10          produces an RNA transcript that interferes with translation of the mRNA of the native DNA sequence.

            Currently, only a few promoters that exhibit a constitutive pattern of expression in plants are available, examples of which include the CaMV 35S, nopaline synthase, and the ubiquitin promoters. The increasing interest in  
15          cotransforming plants with multiple plant transcription units (PTU) and the potential problems associated with using common regulatory sequences for these purposes merit having a number of such promoter sequences available.

            Thus, isolation and characterization of constitutive promoters that can serve as regulatory regions for expression of heterologous nucleotide sequences of  
20          interest are needed for genetic manipulation of plants.

## SUMMARY OF THE INVENTION

            Compositions and methods for regulating expression of heterologous nucleotide sequences in a plant are provided. Compositions are novel nucleotide  
25          sequences for promoters that initiate transcription in a tissue-independent or constitutive manner, more particularly transcriptional initiation regions isolated from the plant genes Actin-2, Enolase, Gos-2 and L41. A method for expressing a heterologous nucleotide sequence in a plant using the transcriptional initiation sequences disclosed herein is provided. The method comprises transforming a  
30          plant cell with a transformation vector that comprises a heterologous nucleotide sequence operably linked to one of the plant promoters of the present invention and regenerating a stably transformed plant from the transformed plant cell. In this

manner, the promoter sequences are useful for controlling the expression of operably linked coding sequences in a constitutive manner.

Downstream from and under the transcriptional initiation regulation of the promoter will be a sequence of interest that will provide for modification of the phenotype of the plant. Such modification includes modulating the production of an endogenous product, as to amount, relative distribution, or the like, or production of an exogenous expression product to provide for a novel function or product in the plant.

## BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the plasmid vector PHP3953 comprising the GUS gene operably linked to the ubiquitin promoter. Promoter fragments of the present invention were recloned into this plasmid in place of the ubiquitin promoter (UBI1ZM), with or without the ubiquitin 5' untranslated region (EX1UBI1ZM) and intron sequence (INT1UBI1ZM), and the resulting plasmid DNA was available in transformation studies to test promoter activity.

Figure 2 schematically depicts an expression cassette for transformation, comprising a gene of interest operably linked to a promoter of the invention.

Figure 3 schematically depicts a plasmid construct comprising the actin-2 promoter of the invention.

Figure 4 schematically depicts a plasmid construct comprising the enolase promoter of the invention.

Figure 5 schematically depicts a plasmid construct comprising the Gos-2 promoter of the invention.

Figure 6 schematically depicts a plasmid construct comprising the L-41 promoter of the invention.

## DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, nucleic acid constructs are provided that allow initiation of transcription in a "tissue-independent," "tissue-general," or "constitutive" manner. Constructs of the invention comprise regulated transcription initiation regions associated with protein translation elongation. Thus, the compositions of the present invention are drawn to novel nucleotide

sequences for tissue-independent, tissue-general, or constitutive plant promoters. By "tissue-independent," "tissue-general," or "constitutive" is intended expression in the cells throughout a plant at most times and in most tissues. As with other promoters classified as "constitutive" (e.g. ubiquitin), some variation in absolute  
5 levels of expression can exist among different tissues or stages.

More particularly, the present invention provides for isolated nucleic acid molecules comprising nucleotide sequences for promoters of the genes actin-2, enolase, Gos-2 and L41. Actins are cytoskeletal proteins which are ubiquitous to nearly all eukaryotes. In plants, several forms of actin are expressed in a variety of  
10 tissues. SEQ ID NO:1 sets forth the nucleotide sequence for the promoter of one of these forms of actin which is present in maize. Enolase is an enzyme of the glycolytic pathway which catalyses the formation of phosphoenolpyruvate from 2-phosphoglycerate. SEQ ID NO:2 sets forth the sequence of a maize enolase promoter.

15 SEQ ID NO: 3 sets forth the sequence for the promoter of GOS-2. SEQ ID NO:4 sets forth the sequence for the promoter of L41. Based on BLAST searches of Genbank databases, L41 cDNA appears to encode a ribosomal protein. Northern blots of root, stem, leaf, ear shoot, kernel and tassie tissues showed mRNA from these genes expressed in many of the tissues.

20 The present invention provides for isolated nucleic acid molecules comprising the nucleotide sequences set forth in SEQ ID NOs: 1-4, or the nucleotide sequences encoding the DNA sequences deposited in a bacterial host as Patent Deposit Nos: 203367, 203368, 203366, and 203369; and fragments and variants thereof

25 Plasmid deposits comprising the promoters of the invention were made to the Patent Depository of American Type Culture Collection (ATCC) on October 21, 1998, and assigned the Patent Deposit No: 203367 for plasmid PHP 13077 comprising the Actin-2 promoter, 203368 for plasmid PHP 13082 comprising the enolase promoter, 203366 for plasmid PHP 13080 comprising the Gos-2 promoter,  
30 and 203369 for plasmid PHP 13084 comprising the L41 promoter. Figures 3, 4, 5 and 6 schematically depict these plasmids. These deposits will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the purposes of Patent Procedure. These deposits

were made merely as a convenience to those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112.

The invention encompasses isolated or substantially purified nucleic acid compositions. An "isolated" or "purified" nucleic acid molecule, or biologically  
5 active portion thereof, is substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. Preferably, an "isolated" nucleic acid is free of sequences (preferably protein encoding sequences) that naturally flank the nucleic acid (i.e., sequences located at the 5' and  
10 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequences that naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived.

15 The promoter nucleotide sequences and methods disclosed herein are useful in regulating constitutive expression of any heterologous nucleotide sequence in a host plant in order to vary the phenotype of a plant.

Various changes in phenotype are of interest including modifying the fatty acid composition in a plant, altering the amino acid content of a plant, altering a  
20 plant's pathogen defense mechanism, and the like. These results can be achieved by providing expression of heterologous products or increased expression of endogenous products in plants. Alternatively, the results can be achieved by providing for a reduction of expression of one or more endogenous products, particularly enzymes or cofactors in the plant. These changes result in a change in  
25 phenotype of the transformed plant.

Genes of interest are reflective of the commercial markets and interests of those involved in the development of the crop. Crops and markets of interest change, and as developing nations open up world markets, new crops and technologies will emerge also. In addition, as our understanding of agronomic  
30 traits and characteristics such as yield and heterosis increase, the choice of genes for transformation will change accordingly. General categories of genes of interest include, for example, those genes involved in information, such as zinc fingers, those involved in communication, such as kinases, and those involved in

housekeeping, such as heat shock proteins. More specific categories of transgenes, for example, include genes encoding important traits for agronomics, insect resistance, disease resistance, herbicide resistance, sterility, grain characteristics, and commercial products. Genes of interest include, generally, those involved in oil, starch, carbohydrate, or nutrient metabolism as well as those affecting kernel size, sucrose loading, and the like.

Agronomically important traits such as oil, starch, and protein content can be genetically altered in addition to using traditional breeding methods. Modifications include increasing content of oleic acid, saturated and unsaturated oils, increasing levels of lysine and sulfur, providing essential amino acids, and also modification of starch. Hordothionin protein modifications are described in U.S. Application Serial No. 08/838,763, filed April 10, 1997; and U.S. Patent Nos. 5,703,049, 5,885,801, and 5,885,802, herein incorporated by reference. Another example is lysine and/or sulfur rich seed protein encoded by the soybean 2S albumin described in U.S. Patent No. 5,850,016, and the chymotrypsin inhibitor from barley, described in Williamson *et al.* (1987) *Eur. J. Biochem.* 165:99-106, the disclosures of which are herein incorporated by reference.

Derivatives of the coding sequences can be made by site-directed mutagenesis to increase the level of preselected amino acids in the encoded polypeptide. For example, the gene encoding the barley high lysine polypeptide (BHL) is derived from barley chymotrypsin inhibitor, U.S. Application Serial No. 08/740,682, filed November 1, 1996, and WO 98/20133, the disclosures of which are herein incorporated by reference. Other proteins include methionine-rich plant proteins such as from sunflower seed (Lilley *et al.* (1989) *Proceedings of the World Congress on Vegetable Protein Utilization in Human Foods and Animal Feedstuffs*, ed. Applewhite (American Oil Chemists Society, Champaign, Illinois), pp. 497-502; herein incorporated by reference); corn (Pedersen *et al.* (1986) *J. Biol. Chem.* 261:6279; Kiriara *et al.* (1988) *Gene* 71:359; both of which are herein incorporated by reference); and rice (Musumura *et al.* (1989) *Plant Mol. Biol.* 12:123, herein incorporated by reference). Other agronomically important genes encode latex, Floury 2, growth factors, seed storage factors, and transcription factors.



Insect resistance genes may encode resistance to pests that have great yield drag such as rootworm, cutworm, European Corn Borer, and the like. Such genes include, for example, *Bacillus thuringiensis* toxic protein genes (U.S. Patent Nos. 5,366,892; 5,747,450; 5,737,514; 5,723,756; 5,593,881; and Geiser *et al.* (1986) *Gene* 48:109); lectins (Van Damme *et al.* (1994) *Plant Mol. Biol.* 24:825); and the like.

Genes encoding disease resistance traits include detoxification genes, such as against fumonisin (U.S. Patent No. 5,792,931); avirulence (avr) and disease resistance (R) genes (Jones *et al.* (1994) *Science* 266:789; Martin *et al.* (1993) *Science* 262:1432; and Mindrinos *et al.* (1994) *Cell* 78:1089); and the like.

Herbicide resistance traits may include genes coding for resistance to herbicides that act to inhibit the action of acetolactate synthase (ALS), in particular the sulfonylurea-type herbicides (e.g., the acetolactate synthase (ALS) gene containing mutations leading to such resistance, in particular the S4 and/or Hra mutations), genes coding for resistance to herbicides that act to inhibit action of glutamine synthase, such as phosphinothricin or basta (e.g., the bar gene), or other such genes known in the art. The *bar* gene encodes resistance to the herbicide basta, the *nptII* gene encodes resistance to the antibiotics kanamycin and geneticin, and the ALS-gene mutants encode resistance to the herbicide chlorsulfuron.

Sterility genes can also be encoded in an expression cassette and provide an alternative to physical detasseling. Examples of genes used in such ways include male tissue-preferred genes and genes with male sterility phenotypes such as QM, described in U.S. Patent No. 5,583,210. Other genes include kinases and those encoding compounds toxic to either male or female gametophytic development.

The quality of grain is reflected in traits such as levels and types of oils, saturated and unsaturated, quality and quantity of essential amino acids, and levels of cellulose. In corn, modified hordothionin proteins are described in copending U.S. Application Serial No. 08/838,763, filed April 10, 1997, and U.S. Patent Nos. 5,703,049, 5,885,801, and 5,885,802.

Commercial traits can also be encoded on a gene or genes that could increase for example, starch for ethanol production, or provide expression of proteins. Another important commercial use of transformed plants is the production of polymers and bioplastics such as described in U.S. Patent No.

5,602,321. Genes such as  $\beta$ -Ketothiolase, PHBase (polyhydroxybutyrate synthase), and acetoacetyl-CoA reductase (see Schubert *et al.* (1988) *J. Bacteriol.* 170:5837-5847) facilitate expression of polyhydroxyalkanoates (PHAs).

Exogenous products include plant enzymes and products as well as those  
5 from other sources including procaryotes and other eukaryotes. Such products include enzymes, cofactors, hormones, and the like. The level of proteins, particularly modified proteins having improved amino acid distribution to improve the nutrient value of the plant, can be increased. This is achieved by the expression of such proteins having enhanced amino acid content.

10 By "promoter" or "transcriptional initiation region" is intended a regulatory region of DNA usually comprising a TATA box capable of directing RNA polymerase II to initiate RNA synthesis at the appropriate transcription initiation site for a particular coding sequence. A promoter may additionally comprise other recognition sequences generally positioned upstream or 5' to the TATA box,  
15 referred to as upstream promoter elements, which influence the transcription initiation rate. It is recognized that having identified the nucleotide sequences for the promoter regions disclosed herein, it is within the state of the art to isolate and identify further regulatory elements in the 5' untranslated region upstream from the particular promoter regions identified herein. Thus the promoter regions disclosed  
20 herein are generally further defined by comprising upstream regulatory elements such as enhancers and the like.

The regulatory sequences of the present invention, when operably linked to a heterologous nucleotide sequence of interest and inserted into a transformation vector, enable constitutive expression of the heterologous nucleotide sequence  
25 throughout the tissues of a plant stably transformed with this vector. That is the nucleotide sequence is expressed at most times and in most tissue types irrespective of developmental stage, environmental stress, and the like. By "heterologous nucleotide sequence" is intended a sequence that is not naturally occurring with the promoter sequence. While this nucleotide sequence is  
30 heterologous to the promoter sequence, it may be homologous, or native, or heterologous, or foreign, to the plant host.

It is recognized that the promoter sequences of the invention may be used to increase or decrease expression of nucleotide sequences that are native or

homologous with respect to the promoter sequence. In this manner, the phenotype of the plant expressing the homologous sequences is altered.

It is also recognized that the nucleotide sequence operably linked to one of the promoter sequences disclosed herein may be an antisense sequence for a particular targeted gene of interest. Antisense constructions are complementary to at least a portion of the messenger RNA (mRNA) for the particular targeted gene, and are constructed to hybridize with the corresponding mRNA. Modifications of the antisense sequences may be made as long as the sequences hybridize to and interfere with expression of the corresponding mRNA. In this manner, antisense constructions having 70%, preferably 80%, more preferably 85% sequence identity to the corresponding antisense sequences may be used. Furthermore, portions of the antisense nucleotides may be used to disrupt the expression of the target gene. Generally, sequences of at least 50 nucleotides, 100 nucleotides, 200 nucleotides, or greater may be used.

The nucleotide sequences of the present invention may also be used with the particular targeted gene sequences in the sense orientation to suppress the expression of endogenous genes in plants. Methods for suppressing gene expression in plants using nucleotide sequences in the sense orientation are known in the art. The methods generally involve transforming plants with a DNA construct comprising a promoter that drives expression in a plant operably linked to at least a portion of a nucleotide sequence that corresponds to the transcript of the endogenous gene. Typically, such a nucleotide sequence has substantial sequence identity to the sequence of the transcript of the endogenous gene, preferably greater than about 65% sequence identity, more preferably greater than about 85% sequence identity, most preferably greater than about 95% sequence identity. See, U.S. Patent Nos. 5,283,184 and 5,034,323; herein incorporated by reference.

The isolated promoter sequences of the present invention can be modified to provide for a range of constitutive expression levels of the heterologous nucleotide sequence. Thus, less than the entire promoter regions may be utilized and the ability to drive expression of the coding sequence retained. However, it is recognized that expression levels of the mRNA may be decreased with deletions of portions of the promoter sequences. Likewise, the tissue-general, constitutive

nature of expression may be changed. Generally, at least about 20 nucleotides of an isolated promoter sequence will be used to drive expression of a nucleotide sequence.

5 It is recognized that to increase transcription levels, enhancers may be utilized in combination with the promoter regions of the invention. Enhancers are nucleotide sequences that act to increase the expression of a promoter region. Enhancers are known in the art and include the SV40 enhancer region, the 35S enhancer element, and the like.

10 Modifications of the isolated promoter sequences of the present invention can provide for a range of constitutive expression of the heterologous nucleotide sequence. Thus, they may be modified to be weak constitutive promoters or strong constitutive promoters. Generally, by "weak promoter" is intended a promoter that drives expression of a coding sequence at a low level. By "low level" is intended at levels of about 1/10,000 transcripts to about 1/100,000 transcripts to about  
15 1/500,000 transcripts. Conversely, a strong promoter drives expression of a coding sequence at a high level, or at about 1/10 transcripts to about 1/100 transcripts to about 1/1,000 transcripts.

Fragments and variants of the disclosed nucleotide sequences are also encompassed by the present invention. By "fragment" is intended a portion of the  
20 nucleotide sequence. Fragments of a promoter nucleotide sequence may retain the biological activity and hence initiate and/or regulate expression. Alternatively, fragments of a nucleotide sequence that are useful as hybridization probes generally do not retain biological activity. Thus, fragments of a nucleotide sequence may range from at least about 20 nucleotides, about 50 nucleotides, about  
25 100 nucleotides, and up to the full-length nucleotide sequence of the invention.

Thus, a fragment of the actin-2, enolase, Gos-2 and L41 promoter nucleotide sequences may encode a biologically active portion or it may be a fragment that can be used as a hybridization probe or PCR primer using methods disclosed below. A biologically active portion can be prepared by isolating a  
30 portion of one of the nucleotide sequences of the invention, and assessing the promoter activity of the portion. Nucleic acid molecules that are fragments of an actin-2, enolase, gos-2 and L41 promoter nucleotide sequence comprise at least 17, 20, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450 nucleotides, or up to the number

of nucleotides present in a full-length nucleotide sequence disclosed herein (for example, 613, 664, 862, and 461 nucleotides for SEQ ID NOs: 1, 2, 3, and 4 respectively).

By "variants" is intended substantially similar sequences. For nucleotide  
5 sequences, variants can be identified with the use of well-known molecular biology techniques, as, for example, with polymerase chain reaction (PCR) and hybridization techniques as outlined below. Variant nucleotide sequences also include synthetically derived nucleotide sequences, such as those generated, for example, by using site-directed mutagenesis. Generally, variants of a particular  
10 nucleotide sequence of the invention will have at least 40%, 50%, 60%, 70%, generally at least 75%, 80%, 85%, preferably about 90% to 95% or more, and more preferably about 98% or more sequence identity to that particular nucleotide sequence as determined by sequence alignment programs described elsewhere herein using default parameters.

15 Variant nucleotide sequences also encompass sequences derived from a mutagenic and recombinogenic procedure such as DNA shuffling. With such a procedure, one or more of the promoter sequences disclosed herein can be manipulated to create a new promoter sequence possessing the desired properties. In this manner, libraries of recombinant polynucleotides are generated from a  
20 population of related sequence polynucleotides comprising sequence regions that have substantial sequence identity and can be homologously recombined *in vitro* or *in vivo*. Strategies for such DNA shuffling are known in the art. See, for example, Stemmer (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751; Stemmer (1994) *Nature* 370:389-391; Crameri *et al.* (1997) *Nature Biotech.* 15:436-438; Moore *et al.* (1997) *J. Mol. Biol.* 272:336-347; Zhang *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:4504-4509; Crameri *et al.* (1998) *Nature* 391:288-291; and U.S. Patent  
25 Nos. 5,605,793 and 5,837,458.

The nucleotide sequences of the invention can be used to isolate corresponding sequences from other organisms, particularly other plants, more  
30 particularly other monocots. In this manner, methods such as PCR, hybridization, and the like can be used to identify such sequences based on their sequence homology to the sequences set forth herein. Sequences isolated based on their

sequence identity to the entire actin-2, enolase, Gos-2 or L41 promoter sequences set forth herein, or to fragments thereof are encompassed by the present invention.

In a PCR approach, oligonucleotide primers can be designed for use in PCR reactions to amplify corresponding DNA sequences from cDNA or genomic DNA extracted from any plant of interest. Methods for designing PCR primers and PCR cloning are generally known in the art and are disclosed in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York). See also Innis *et al.*, eds. (1990) *PCR Protocols: A Guide to Methods and Applications* (Academic Press, New York); Innis and Gelfand, eds. (1995) *PCR Strategies* (Academic Press, New York); and Innis and Gelfand, eds. (1999) *PCR Methods Manual* (Academic Press, New York). Known methods of PCR include, but are not limited to, methods using paired primers, nested primers, single specific primers, degenerate primers, gene-specific primers, vector-specific primers, partially-mismatched primers, and the like.

In hybridization techniques, all or part of a known nucleotide sequence is used as a probe that selectively hybridizes to other corresponding nucleotide sequences present in a population of cloned genomic DNA fragments or cDNA fragments (i.e., genomic or cDNA libraries) from a chosen organism. The hybridization probes may be genomic DNA fragments, cDNA fragments, RNA fragments, or other oligonucleotides, and may be labeled with a detectable group such as  $^{32}\text{P}$ , or any other detectable marker. Thus, for example, probes for hybridization can be made by labeling synthetic oligonucleotides based on the promoter sequences of the invention. Methods for preparation of probes for hybridization and for construction of cDNA and genomic libraries are generally known in the art and are disclosed in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

For example, the entire actin-2, enolase, Gos-2 or L41 promoter sequence disclosed herein, or one or more portions thereof, may be used as a probe capable of specifically hybridizing to corresponding actin-2, enolase, Gos-2 or L41 sequences. To achieve specific hybridization under a variety of conditions, such probes include sequences that are unique among e promoter sequences and are preferably at least about 10 nucleotides in length, and most preferably at least

about 20 nucleotides in length. Such probes may be used to amplify corresponding plant sequences from a chosen plant by PCR. This technique may be used to isolate additional coding sequences from a desired plant or as a diagnostic assay to determine the presence of coding sequences in a plant. Hybridization techniques include hybridization screening of plated DNA libraries (either plaques or colonies; see, for example, Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

Hybridization of such sequences may be carried out under stringent conditions. By "stringent conditions" or "stringent hybridization conditions" is intended conditions under which a probe will hybridize to its target sequence to a detectably greater degree than to other sequences (e.g., at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences that are 100% complementary to the probe can be identified (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, preferably less than 500 nucleotides in length.

Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C, and a wash in 1X to 2X SSC (20X SSC = 3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55°C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1.0 M NaCl, 1% SDS at 37°C, and a wash in 0.5X to 1X SSC at 55 to 60°C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.1X SSC at 60 to 65°C.

Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the  $T_m$  can be approximated from the equation of Meinkoth and Wahl (1984) *Anal. Biochem.* 138:267-284:  $T_m = 81.5^\circ\text{C} + 16.6 (\log M) + 0.41 (\%GC) - 0.61 (\% \text{ form}) - 500/L$ ; where M is the molarity of monovalent cations, %GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe.  $T_m$  is reduced by about  $1^\circ\text{C}$  for each 1% of mismatching; thus,  $T_m$ , hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with  $\geq 90\%$  identity are sought, the  $T_m$  can be decreased  $10^\circ\text{C}$ . Generally, stringent conditions are selected to be about  $5^\circ\text{C}$  lower than the thermal melting point ( $T_m$ ) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or  $4^\circ\text{C}$  lower than the thermal melting point ( $T_m$ ); moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or  $10^\circ\text{C}$  lower than the thermal melting point ( $T_m$ ); low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or  $20^\circ\text{C}$  lower than the thermal melting point ( $T_m$ ). Using the equation, hybridization and wash compositions, and desired  $T_m$ , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a  $T_m$  of less than  $45^\circ\text{C}$  (aqueous solution) or  $32^\circ\text{C}$  (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 (Elsevier, New York); and Ausubel *et al.*, eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2 (Greene Publishing and Wiley-Interscience, New York). See Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York).



Thus, isolated sequences that have promoter activity and which hybridize under stringent conditions to the promoter sequences disclosed herein, or to fragments thereof, are encompassed by the present invention. Such sequences will be at least 40% to 50% homologous, about 60% to 70% homologous, and even  
5 about 75%, 80%, 85%, 90%, 95% to 98% homologous or more with the disclosed sequences. That is, the sequence identity of sequences may range, sharing at least 40% to 50%, about 60% to 70%, and even about 75%, 80%, 85%, 90%, 95% to 98% or more sequence identity. The promoter regions of the invention may be used to isolate substantially identical sequences from any plant species, including  
10 but not limited to any plant species described herein.

The following terms are used to describe the sequence relationships between two or more nucleic acids or polynucleotides: (a) "reference sequence", (b) "comparison window", (c) "sequence identity", (d) "percentage of sequence identity", and (e) "substantial identity".

15 (a) As used herein, "reference sequence" is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full-length cDNA or gene sequence, or the complete cDNA or gene sequence.

(b) As used herein, "comparison window" makes reference to a  
20 contiguous and specified segment of a polynucleotide sequence, wherein the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Generally, the comparison window is at least 20 contiguous nucleotides in length, and optionally  
25 can be 30, 40, 50, 100, or longer. Those of skill in the art understand that to avoid a high similarity to a reference sequence due to inclusion of gaps in the polynucleotide sequence a gap penalty is typically introduced and is subtracted from the number of matches.

Methods of alignment of sequences for comparison are well known in the  
30 art. Thus, the determination of percent identity between any two sequences can be accomplished using a mathematical algorithm. Preferred, non-limiting examples of such mathematical algorithms are the algorithm of Myers and Miller (1988) *CABIOS* 4:11-17; the local homology algorithm of Smith *et al.* (1981) *Adv. Appl.*

- Math.* 2:482; the homology alignment algorithm of Needleman *and Wunsch* (1970) *J. Mol. Biol.* 48:443-453; the search-for-similarity-method of Pearson *and Lipman* (1988) *Proc. Natl. Acad. Sci.* 85:2444-2448; the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264, modified as in Karlin and Altschul
- 5 (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877.

Computer implementations of these mathematical algorithms can be utilized for comparison of sequences to determine sequence identity. Such implementations include, but are not limited to: CLUSTAL in the PC/Gene program (available from Intelligenetics, Mountain View, California); the ALIGN

10 program (Version 2.0) and GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Version 8 (available from Genetics Computer Group (GCG), 575 Science Drive, Madison, Wisconsin, USA). Alignments using these programs can be performed using the default parameters. The CLUSTAL program is well described by Higgins *et al.* (1988) *Gene* 73:237-

15 244 (1988); Higgins *et al.* (1989) *CABIOS* 5:151-153; Corpet *et al.* (1988) *Nucleic Acids Res.* 16:10881-90; Huang *et al.* (1992) *CABIOS* 8:155-65; and Pearson *et al.* (1994) *Meth. Mol. Biol.* 24:307-331. The ALIGN program is based on the algorithm of Myers and Miller (1988) *supra*. The BLAST programs of Altschul *et al.* (1990) *J. Mol. Biol.* 215:403 are based on the algorithm of Karlin and Altschul

20 (1990) *supra*. To obtain gapped alignments for comparison purposes, Gapped BLAST (in BLAST 2.0) can be utilized as described in Altschul *et al.* (1997) *Nucleic Acids Res.* 25:3389. Alternatively, PSI-BLAST (in BLAST 2.0) can be used to perform an iterated search that detects distant relationships between molecules. See Altschul *et al.* (1997) *supra*. When utilizing BLAST, Gapped

25 BLAST, PSI-BLAST, the default parameters of the respective programs (e.g., BLASTN for nucleotide sequences, BLASTX for proteins) can be used. See <http://www.ncbi.nlm.nih.gov>. Alignment may also be performed manually by inspection. Alignment may also be performed manually by inspection.

For purposes of the present invention, comparison of nucleotide sequences

30 for determination of percent sequence identity to the promoter sequences disclosed herein is preferably made using the BlastN program (version 1.4.7 or later) with its default parameters or any equivalent program. By "equivalent program" is intended any sequence comparison program that, for any two sequences in

question, generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by the preferred program.

(c) As used herein, "sequence identity" or "identity" in the context of two nucleic acid or polypeptide sequences makes reference to the residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to proteins it is recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have "sequence similarity" or "similarity". Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, California).

(d) As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity.

(e)(i) The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 70% sequence identity, preferably at least 80%, more preferably at least 90%, and most preferably at least 95%, compared to a reference sequence using one of the alignment programs described using standard parameters. One of skill in the art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning, and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 60%, more preferably at least 70%, 80%, 90%, and most preferably at least 95%.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other under stringent conditions. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. However, stringent conditions encompass temperatures in the range of about 1°C to about 20°C, depending upon the desired degree of stringency as otherwise qualified herein. Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides they encode are substantially identical. This may occur, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. One indication that two nucleic acid sequences are substantially identical is when the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid.

(e)(ii) The term "substantial identity" in the context of a peptide indicates that a peptide comprises a sequence with at least 70% sequence identity to a reference sequence, preferably 80%, more preferably 85%, most preferably at least 90% or 95% sequence identity to the reference sequence over a specified comparison window. Preferably, optimal alignment is conducted using the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443-453. An indication that two peptide sequences are substantially identical is that one peptide is immunologically reactive with antibodies raised against the second peptide. Thus, a peptide is substantially identical to a second peptide, for

example, where the two peptides differ only by a conservative substitution. Peptides that are "substantially similar" share sequences as noted above except that residue positions that are not identical may differ by conservative amino acid changes.

5           The nucleotide sequences for the constitutive promoters disclosed in the present invention, as well as variants and fragments thereof, are useful in the genetic manipulation of any plant when operably linked with a nucleotide sequence whose constitutive expression is to be controlled to achieve a desired phenotypic response. By "operably linked" is intended the transcription or translation of the  
10       heterologous nucleotide sequence is under the influence of the promoter sequence. In this manner, the nucleotide sequences for the promoters of the invention may be provided in expression cassettes along with nucleotide sequences of interest for expression in the plant of interest.

          Such expression cassettes will comprise a transcriptional initiation region  
15       comprising one of the promoter nucleotide sequences of the present invention, or variants or fragments thereof, operably linked to the nucleotide sequence whose expression is to be controlled by the constitutive promoters disclosed herein. Such an expression cassette is provided with a plurality of restriction sites for insertion of the nucleotide sequence to be under the transcriptional regulation of the  
20       regulatory regions. The expression cassette may additionally contain selectable marker genes.

          The transcriptional cassette will include in the 5'- to -3' direction of transcription, a transcriptional and translational initiation region, a heterologous nucleotide sequence of interest, and a transcriptional and translational termination  
25       region functional in plants. The termination region may be native with the transcriptional initiation region comprising one of the promoter nucleotide sequences of the present invention, may be native with the DNA sequence of interest, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase  
30       and nopaline synthase termination regions. See also, Guerineau *et al.* (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon *et al.* (1991) *Genes Dev.* 5:141-149; Mogen *et al.* (1990) *Plant Cell* 2:1261-1272;

Munroe *et al.* (1990) *Gene* 91:151-158; Ballas *et al.* (1989) *Nucleic Acids Res.* 17:7891-7903; Joshi heterologous *et al.* (1987) *Nucleic Acid Res.* 15:9627-9639.

The expression cassette comprising the promoter sequence of the present invention operably linked to a nucleotide sequence may also contain at least one  
5 additional nucleotide sequence for a gene to be cotransformed into the organism. Alternatively, the additional sequence(s) can be provided on another expression cassette.

Where appropriate, the heterologous nucleotide sequence whose expression is to be under the control of the promoter sequence of the present invention and  
10 any additional nucleotide sequence(s) may be optimized for increased expression in the transformed plant. That is, these nucleotide sequences can be synthesized using plant preferred codons for improved expression. Methods are available in the art for synthesizing plant-preferred nucleotide sequences. See, for example, U.S. Patent Nos. 5,380,831 and 5,436,391, and Murray *et al.* (1989) *Nucleic Acids*  
15 *Res.* 17:477-498, herein incorporated by reference.

Additional sequence modifications are known to enhance gene expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon-intron splice site signals, transposon-like repeats, and other such well-characterized sequences that may be deleterious to gene  
20 expression. The G-C content of the heterologous nucleotide sequence may be adjusted to levels average for a given cellular host, as calculated by reference to known genes expressed in the host cell. When possible, the sequence is modified to avoid predicted hairpin secondary mRNA structures.

The expression cassettes may additionally contain 5' leader sequences in  
25 the expression cassette construct. Such leader sequences can act to enhance translation. Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5' noncoding region) (Elroy-Stein *et al.* (1989) *Proc. Nat. Acad. Sci. USA* 86:6126-6130); potyvirus leaders, for example, TEV leader (Tobacco Etch Virus) (Allison *et al.* (1986));  
30 MDMV leader (Maize Dwarf Mosaic Virus) (*Virology* 154:9-20); human immunoglobulin heavy-chain binding protein (BiP) (Macejak *et al.* (1991) *Nature* 353:90-94); untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4) (Jobling *et al.* (1987) *Nature* 325:622-625); tobacco mosaic

virus leader (TMV) (Gallie *et al.* (1989) *Molecular Biology of RNA*, pages 237-256); and maize chlorotic mottle virus leader (MCMV) (Lommel *et al.* (1991) *Virology* 81:382-385). See also Della-Cioppa *et al.* (1987) *Plant Physiology* 84:965-968. Other methods known to enhance translation and/or mRNA stability  
5 can also be utilized, for example, introns, and the like.

In those instances where it is desirable to have the constitutively expressed product of the heterologous nucleotide sequence directed to a particular organelle, such as the chloroplast or mitochondrion, or secreted at the cell's surface or extracellularly, the expression cassette may further comprise a coding sequence for  
10 a transit peptide. Such transit peptides are well known in the art and include, but are not limited to, the transit peptide for the acyl carrier protein, the small subunit of RUBISCO, plant EPSP synthase, and the like.

In preparing the expression cassette, the various DNA fragments may be manipulated, so as to provide for the DNA sequences in the proper orientation and,  
15 as appropriate, in the proper reading frame. Toward this end, adapters or linkers may be employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites, or the like. For this purpose, *in vitro* mutagenesis, primer repair, restriction, annealing, resubstitutions, for example, transitions and  
20 transversions, may be involved.

The promoters may be used to drive reporter genes and/or selectable marker genes. Examples of suitable reporter genes known in the art can be found in, for example, Jefferson *et al.* (1991) in *Plant Molecular Biology Manual*, ed. Gelvin *et al.* (Kluwer Academic Publishers), pp. 1-33; DeWet *et al.* (1987) *Mol.*  
25 *Cell. Biol.* 7:725-737; Goff *et al.* (1990) *EMBO J.* 9:2517-2522; Kain *et al.* (1995) *BioTechniques* 19:650-655; and Chiu *et al.* (1996) *Current Biology* 6:325-330.

Selectable marker genes for selection of transformed cells or tissues can include genes that confer antibiotic resistance or resistance to herbicides. Examples of suitable selectable marker genes include, but are not limited to, genes  
30 encoding resistance to chloramphenicol (Herrera Estrella *et al.* (1983) *EMBO J.* 2:987-992); methotrexate (Herrera Estrella *et al.* (1983) *Nature* 303:209-213; Meijer *et al.* (1991) *Plant Mol. Biol.* 16:807-820); hygromycin (Waldron *et al.* (1985) *Plant Mol. Biol.* 5:103-108; Zhijian *et al.* (1995) *Plant Science* 108:219-

227); streptomycin (Jones *et al.* (1987) *Mol. Gen. Genet.* 210:86-91); spectinomycin (Bretagne-Sagnard *et al.* (1996) *Transgenic Res.* 5:131-137); bleomycin (Hille *et al.* (1990) *Plant Mol. Biol.* 7:171-176); sulfonamide (Guerineau *et al.* (1990) *Plant Mol. Biol.* 15:127-136); bromoxynil (Stalker *et al.* (1988) *Science* 242:419-423); glyphosate (Shaw *et al.* (1986) *Science* 233:478-481); phosphinothricin (DeBlock *et al.* (1987) *EMBO J.* 6:2513-2518).

Other genes that could serve utility in the recovery of transgenic events but might not be required in the final product would include, but are not limited to, examples such as GUS (b-glucuronidase; Jefferson (1987) *Plant Mol. Biol. Rep.* 5:387), GFP (green fluorescence protein; Chalfie *et al.* (1994) *Science* 263:802), luciferase (Riggs *et al.* (1987) *Nucleic Acid Res.* 15(19):8115; Luehrsen *et al.* (1992) *Methods Enzymol.* 216:397-414) and the maize genes encoding for anthocyanin production (Ludwig *et al.* (1990) *Science* 247:449).

The expression cassette comprising the particular promoter sequence of the present invention operably linked to a nucleotide sequence of interest can be used to transform any plant. In this manner, genetically modified plants, plant cells, plant tissue, seed, and the like can be obtained. Transformation protocols as well as protocols for introducing nucleotide sequences into plants may vary depending on the type of plant or plant cell, i.e., monocot or dicot, targeted for transformation. Suitable methods of introducing nucleotide sequences into plant cells and subsequent insertion into the plant genome include microinjection (Crossway *et al.* (1986) *Biotechniques* 4:320-334), electroporation (Riggs *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606, *Agrobacterium*-mediated transformation (Townsend *et al.*, U.S. Pat No. 5,563,055), direct gene transfer (Paszkowski *et al.* (1984) *EMBO J.* 3:2717-2722), and ballistic particle acceleration (see, for example, Sanford *et al.*, U.S. Patent No. 4,945,050; Tomes *et al.*, U.S. Patent No. 5,879,918; Tomes *et al.*, U.S. Patent No. 5,886,244; Bidney *et al.*, U.S. Patent No. 5,932,782; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); and McCabe *et al.* (1988) *Biotechnology* 6:923-926). Also see Weissinger *et al.* (1988) *Ann. Rev. Genet.* 22:421-477; Sanford *et al.* (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou *et al.* (1988) *Plant Physiol.* 87:671-674



(soybean); McCabe *et al.* (1988) *Bio/Technology* 6:923-926 (soybean); Finer and McMullen (1991) *In Vitro Cell Dev. Biol.* 27P:175-182 (soybean); Singh *et al.* (1998) *Theor. Appl. Genet.* 96:319-324 (soybean); Datta *et al.* (1990) *Biotechnology* 8:736-740 (rice); Klein *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize); Klein *et al.* (1988) *Biotechnology* 6:559-563 (maize);  
 5 Tomes, U.S. Patent No. 5,240,855; Buising *et al.*, U.S. Patent Nos. 5,322,783 and 5,324,646; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment," in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg (Springer-Verlag, Berlin) (maize); Klein *et al.* (1988) *Plant Physiol.* 91:440-444 (maize); Fromm *et al.* (1990) *Biotechnology* 8:833-839 (maize); Hooykaas-Van Slogteren *et al.* (1984) *Nature (London)* 311:763-764; Bowen *et al.*, U.S. Patent No. 5,736,369 (cereals); Bytebier *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:5345-5349 (Liliaceae); De Wet *et al.* (1985) in *The Experimental Manipulation of Ovule Tissues*, ed. Chapman *et al.* (Longman, New York), pp. 197-209 (pollen); Kaeppler *et al.* (1990) *Plant Cell Reports* 9:415-418 and Kaeppler *et al.* (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated transformation); D'Halluin *et al.* (1992) *Plant Cell* 4:1495-1505 (electroporation); Li *et al.* (1993) *Plant Cell Reports* 12:250-255 and Christou and Ford (1995) *Annals of Botany* 75:407-413 (rice); Osjoda *et al.* (1996) *Nature Biotechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); all of which are herein  
 10  
 15  
 20 incorporated by reference.

The cells that have been transformed may be grown into plants in accordance with conventional ways. See, for example, McCormick *et al.* (1986) *Plant Cell Reports* 5:81-84. These plants may then be grown, and either pollinated  
 25 with the same transformed strain or different strains, and the resulting hybrid having constitutive expression of the desired phenotypic characteristic identified. Two or more generations may be grown to ensure that expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure expression of the desired phenotypic characteristic has been  
 30 achieved.

The present invention may be used for transformation of any plant species, including, but not limited to, corn (*Zea mays*), *Brassica* sp. (e.g., *B. napus*, *B. rapa*, *B. juncea*), particularly those *Brassica* species useful as sources of seed oil, alfalfa

- (*Medicago sativa*), rice (*Oryza sativa*), rye (*Secale cereale*), sorghum (*Sorghum bicolor*, *Sorghum vulgare*), millet (e.g., pearl millet (*Pennisetum glaucum*), proso millet (*Panicum miliaceum*), foxtail millet (*Setaria italica*), finger millet (*Eleusine coracana*)), sunflower (*Helianthus annuus*), safflower (*Carthamus tinctorius*), wheat
- 5 (*Triticum aestivum*), soybean (*Glycine max*), tobacco (*Nicotiana tabacum*), potato (*Solanum tuberosum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium barbadense*, *Gossypium hirsutum*), sweet potato (*Ipomoea batatas*), cassava (*Manihot esculenta*), coffee (*Cofea* spp.), coconut (*Cocos nucifera*), pineapple (*Ananas comosus*), citrus trees (*Citrus* spp.), cocoa (*Theobroma cacao*), tea (*Camellia sinensis*), banana (*Musa*
- 10 spp.), avocado (*Persea americana*), fig (*Ficus casica*), guava (*Psidium guajava*), mango (*Mangifera indica*), olive (*Olea europaea*), papaya (*Carica papaya*), cashew (*Anacardium occidentale*), macadamia (*Macadamia integrifolia*), almond (*Prunus amygdalus*), sugar beets (*Beta vulgaris*), sugarcane (*Saccharum* spp.), oats, barley, vegetables, ornamentals, and conifers.
- 15 Vegetables include tomatoes (*Lycopersicon esculentum*), lettuce (e.g., *Lactuca sativa*), green beans (*Phaseolus vulgaris*), lima beans (*Phaseolus limensis*), peas (*Lathyrus* spp.), and members of the genus *Cucumis* such as cucumber (*C. sativus*), cantaloupe (*C. cantalupensis*), and musk melon (*C. melo*). Ornamentals include azalea (*Rhododendron* spp.), hydrangea (*Macrophylla*
- 20 *hydrangea*), hibiscus (*Hibiscus rosasanensis*), roses (*Rosa* spp.), tulips (*Tulipa* spp.), daffodils (*Narcissus* spp.), petunias (*Petunia hybrida*), carnation (*Dianthus caryophyllus*), poinsettia (*Euphorbia pulcherrima*), and chrysanthemum. Conifers that may be employed in practicing the present invention include, for example, pines such as loblolly pine (*Pinus taeda*), slash pine (*Pinus elliotii*), ponderosa pine
- 25 (*Pinus ponderosa*), lodgepole pine (*Pinus contorta*), and Monterey pine (*Pinus radiata*); Douglas-fir (*Pseudotsuga menziesii*); Western hemlock (*Tsuga canadensis*); Sitka spruce (*Picea glauca*); redwood (*Sequoia sempervirens*); true firs such as silver fir (*Abies amabilis*) and balsam fir (*Abies balsamea*); and cedars such as Western red cedar (*Thuja plicata*) and Alaska yellow-cedar
- 30 (*Chamaecyparis nootkatensis*). Preferably, plants of the present invention are crop plants (for example, corn, alfalfa, sunflower, *Brassica*, soybean, cotton, safflower, peanut, sorghum, wheat, millet, tobacco, etc.), more preferably corn and soybean plants, yet more preferably corn plants.

Plants of particular interest include grain plants that provide seeds of interest, oil-seed plants, and leguminous plants. Seeds of interest include grain seeds, such as corn, wheat, barley, rice, sorghum, rye, etc. Oil-seed plants include cotton, soybean, safflower, sunflower, *Brassica*, maize, alfalfa, palm, coconut, etc.

- 5 Leguminous plants include beans and peas. Beans include guar, locust bean, fenugreek, soybean, garden beans, cowpea, mungbean, lima bean, fava bean, lentils, chickpea, etc.

The following examples are offered by way of illustration and not by way of limitation.

10

## EXPERIMENTAL

- Promoter regions for the maize genes actin-2, enolase, Gos-2 and L41 were isolated from maize plants and cloned. The sequences for these promoters are set forth in SEQ ID NOs: 1, 2, 3 and 4 respectively. The method for their isolation is described below.
- 15

### Example 1: Isolation and Cloning of the Promoters

- Pioneer-proprietary maize EST (expressed sequence tag) library was examined by computer and manually to identify candidate genes of frequency among the various entries and distribution among maize tissues so as to be likely expressed ubiquitously (or nearly so) in maize and at relatively high mRNA levels. Approximately ten candidate genes were so identified. Northern blots were run with various maize tissue mRNA samples, probed with these ESTs. Those ESTs yielding apparently good mRNA expression were selected for cloning of the corresponding promoter fragments. The putative promoters were isolated from genomic maize DNA according to methods and materials provided in the Genome Walker Kit by Clontech Corporation, and as described in more detail below. Four of these putative promoters were cloned upstream of a GUS reporter cDNA (PHP3953, Figure 1). For each of these promoters, two types of constructs were generated, one with and the other without the maize ubiquitin 5'UT (EX1UBI1ZM) and intron sequence (INT1UBI1ZM of PHP3953).
- 20
- 25
- 30

This plus/minus INT1UBI1ZM sequence was to allow testing for whether that ubi sequence significantly aided the transcriptional activity of the promoters.

The constructs are used to test transient and stable maize transformation, with scoring of GUS expression levels, temporally and developmentally relative to the parent PHP3953 construct which contains the maize Ubiquitin promoter/ intron/ and GUS reporter (Figure 1), and to a basal construct containing only the GUS reporter.

The procedure for promoter isolation is described in the user manual for the GenomeWalker kit sold by Clontech Laboratories, Inc., Palo Alto, CA. Genomic DNA libraries of line B73 were constructed by digesting the maize genomic DNA with 5 different enzymes DraI, Eco RV, PvuII, ScaI and StuI. The resulting DNA is referred to as DL1-DL5, respectively. For isolation of specific promoter regions, two non-overlapping gene-specific primers were designed from the sequences of abundant ESTs in the Pioneer/HGS database. The primers were designed to amplify the region upstream of the coding sequence, i.e. the 5' -untranslated region and promoter region of the chosen gene. The sequences of the primers are given below for each promoter described. The first round of PCR was performed on each DNA sample (DL1-5) with Clontech Primer AP1 (sequence 5'-gtaatacgaactcactatagggc- 3') and the gene-specific primer (gsp1) with the following sequences:

Actin-2	gsp1	:5'- cttggactgggcctcgctaccaacata -3'
Gos-2	gsp1	:5'- accagcaccagagtcctcagcattgcctcag -3'
Enolase	gsp1	:5'- ttgcctcgcaagtcacgctcgattcgg -3'
L41	gsp1	:5'- ttccctgggcagacaggctatccttacc -3'

20

PCR was performed in a 2400 Thermal Cycler with heated lid from Perkin Elmer (Foster City, CA) using reagents supplied with the GenomeWalker kit. The following cycling parameters were used: seven cycles of 94° C for 2 sec., then 72° C for 3 min., followed by 32 cycles of 94° C for 2 sec., 67° C for 3 min. Finally, the samples were held at 67° C for 4 min., then at 4° C until analyzed.

As described in the user manual, the DNA from the first round of PCR was then diluted and used as a template in a second round of PCR using the Clontech AP2 primer (sequence 5'- actatagggcagcggtgt-3') and gene specific primer (gsp)2 with the following sequences:

Actin-2	gsp2:	5'- actgatgatgccggagatctaggagaag-3'
Gos-2	gsp2:	5'- tcgcttctctggaggaggcgtggagcgac-3'
Enolase	gsp2:	5'- ccacttggcgggtgacagatgttcgagc-3'
L41	gsp2:	5'- cttggtcttaggaacgttcaccatttggcgg-3'

The cycling parameters for the second round were: five cycles of 94° C for 2 sec., then 72° C for 3 min., followed by 20 cycles of 94° C for 2 sec., and 67° C for 3 min., finally four min. at 67° C and then held at 4° C. Approximately ten µl of each reaction were run on a 1% agarose gel, and bands larger than 500 bp were excised, purified with Wizard PCR Prep Kit (Promega, Madison, WI) and cloned into the TA vector pCR 2.1 (Invitrogen, San Diego, CA). Clones were sequenced for verification, and then 2µl was amplified with gene specific primer gsp3 (sequence below) and the adapter primer AP3 to add recognition sequences for restriction enzymes to ease the cloning process. This amplification was according to the following cycling parameters: five cycles of 94° C for 2 sec., 46° C for 30 sec., 72° C for 3 min., followed by 20 cycles of 94° C for 2 sec., 67° C for 3 min., finally 67° C for 4 min., and held at 4° C.

15

AP3	:	5'- atatgcatgcatctgcagatcccgggactatagggcacgcgtggt-3'
Acti-2	Gsp3:	5'- atccatggtgtcgtgtggatccactgatgatgccggagatctagg-3'
Gos-2	Gsp3:	5'- atccatggtgtcgtgtggatcctcgttctctggaggaggcgtg-3'
Enolase	Gsp3:	5'- atccatggtgtcgtgtggatccacttggcgggtgacagatgttc-3'
L41	Gsp3:	5'- atccatggtgtcgtgtggatcccttgggtcttaggaacgttcacca-3'

Ten ul of the resulting amplified DNA was run on a 1% agarose gel, purified with Wizard PCR Prep Kit and digested with SphI and BamHI, run on gel, purified, then subcloned into PHP3953 cut with SphI and BglII, thereby substituting the ubiquitin promoter. Final sequences were determined for the resulting plasmids, the pertinent parts of which are set forth in SEQ ID NOs: 1-4. The sequences of these four promoters showed no significant match to other transcriptional promoters when checked by GCG BlastN analyses of the GeneSeqN database. The GeneSeq database is published by Derwent Scientific

Publications and distributed electronically by Oxford Molecular Group. Oxford Molecular coordinates the content assembly of the GeneSeq database with Derwent. The GeneSeq database is compiled for institutional and corporate use by GCG, which is also now a part of Oxford Molecular.

5

#### Example 2: Expression Using Novel Maize Promoters

A transient expression assay is used to test the cloned DNAs for promoter activity. The promoter sequences of SEQ ID NOs:1-4 were cloned into a GUS expression vector (Figure 1) as described above in Example 1.

10

Plasmid DNA was bombarded into maize cell suspension culture and GUS activity is measured, using the ubiquitin promoter as a control, by counting blue spots after staining for GUS activity as described in Jefferson (1987) *Plant Mol. Biol. Rep.* 5:387-405.

15

#### Example 3: Transformation and Regeneration of Transgenic Plants

Immature maize embryos from greenhouse donor plants are bombarded with a plasmid containing an actin-2, enolase, Gos-2 or L41 promoter of the invention operably linked to a gene of interest plus a plasmid containing the selectable marker gene PAT (Wohlleben *et al.* (1988) *Gene* 70:25-37) that confers resistance to the herbicide Bialaphos. Transformation is performed as follows. All media recipes are in the Appendix.

25

#### Preparation of Target Tissue

The ears are surface sterilized in 30% Chlorox bleach plus 0.5% Micro detergent for 20 minutes, and rinsed two times with sterile water. The immature embryos are excised and placed embryo axis side down (scutellum side up), 25 embryos per plate, on 560Y medium for 4 hours and then aligned within the 2.5-cm target zone in preparation for bombardment.

30

### Preparation of DNA

A plasmid vector comprising the gene of interest operably linked to an actin-2, enolase, Gos-2 or L41 promoter of the invention (SEQ ID NOs: 1-4 respectively) is made. The respective constructs comprising these promoters are shown in Figures 3, 4, 5, and 6. This plasmid DNA plus plasmid DNA containing a PAT selectable marker is precipitated onto 1.1  $\mu\text{m}$  (average diameter) tungsten pellets using a  $\text{CaCl}_2$  precipitation procedure as follows:

- 100  $\mu\text{l}$  prepared tungsten particles in water
- 10  $\mu\text{l}$  (1  $\mu\text{g}$ ) DNA in TrisEDTA buffer (1  $\mu\text{g}$  total)
- 100  $\mu\text{l}$  2.5 M  $\text{CaCl}_2$
- 10  $\mu\text{l}$  0.1 M spermidine

- Each reagent is added sequentially to the tungsten particle suspension, while maintained on the multitube vortexer. The final mixture is sonicated briefly and allowed to incubate under constant vortexing for 10 minutes. After the precipitation period, the tubes are centrifuged briefly, liquid removed, washed with 500  $\mu\text{l}$  100% ethanol, and centrifuged for 30 seconds. Again the liquid is removed, and 105  $\mu\text{l}$  100% ethanol is added to the final tungsten particle pellet.
- For particle gun bombardment, the tungsten/DNA particles are briefly sonicated and 10  $\mu\text{l}$  spotted onto the center of each macrocarrier and allowed to dry about 2 minutes before bombardment.

### Particle Gun Treatment

- The sample plates are bombarded at level #4 in particle gun #HE34-1 or #HE34-2. All samples receive a single shot at 650 PSI, with a total of ten aliquots taken from each tube of prepared particles/DNA.

### Subsequent Treatment

- Following bombardment, the embryos are kept on 560Y medium for 2 days, then transferred to 560R selection medium containing 3 mg/liter Bialaphos, and subcultured every 2 weeks. After approximately 10 weeks of selection,

selection-resistant callus clones are transferred to 288J medium to initiate plant regeneration. Following somatic embryo maturation (2-4 weeks), well-developed somatic embryos are transferred to medium for germination and transferred to the lighted culture room. Approximately 7-10 days later, developing plantlets are

5 transferred to 272V hormone-free medium in tubes for 7-10 days until plantlets are well established. Plants are then transferred to inserts in flats (equivalent to 2.5" pot) containing potting soil and grown for 1 week in a growth chamber, subsequently grown an additional 1-2 weeks in the greenhouse, then transferred to classic 600 pots (1.6 gallon) and grown to maturity. Plants are monitored and

10 scored for expression and/or activity of the gene of interest.



**APPENDIX****272 V**

Ingredient	Amount	Unit
D-I H <sub>2</sub> O	950.000	ml
MS Salts (GIBCO 11117-074)	4.300	G
Myo-Inositol	0.100	G
MS Vitamins Stock Solution ##	5.000	ml
Sucrose	40.000	G
Bacto-Agar @	6.000	G

5

**Directions:****@ = Add after bringing up to volume****Dissolve ingredients in polished D-I H<sub>2</sub>O in sequence****Adjust to pH 5.6**10 **Bring up to volume with polished D-I H<sub>2</sub>O after adjusting pH****Sterilize and cool to 60°C.**

**## = Dissolve 0.100 g of Nicotinic Acid; 0.020 g of Thiamine.HCL; 0.100 g of Pyridoxine.HCL; and 0.400 g of Glycine in 875.00 ml of polished D-I H<sub>2</sub>O in sequence. Bring up to volume with polished D-I H<sub>2</sub>O. Make in 400 ml portions.**

15 **Thiamine.HCL & Pyridoxine.HCL are in Dark Desiccator. Store for one month, unless contamination or precipitation occurs, then make fresh stock.**

**Total Volume (L) = 1.00**

288 J

Ingredient	Amount	Unit
D-I H <sub>2</sub> O	950.000	MI
MS Salts	4.300	G
Myo-Inositol	0.100	G
MS Vitamins Stock Solution ##	5.000	MI
Zeatin .5mg/ml	1.000	MI
Sucrose	60.000	G
Gelrite @	3.000	G
Indoleacetic Acid 0.5 mg/ml #	2.000	MI
0.1mM Absciscic Acid	1.000	MI
Bialaphos 1mg/ml #	3.000	MI

## Directions:

@ = Add after bringing up to volume

- 5 Dissolve ingredients in polished D-I H<sub>2</sub>O in sequence

Adjust to pH 5.6

Bring up to volume with polished D-I H<sub>2</sub>O after adjusting pH

Sterilize and cool to 60°C.

Add 3.5g/L of Gelrite for cell biology.

- 10 ## = Dissolve 0.100 g of Nicotinic Acid; 0.020 g of Thiamine.HCL; 0.100 g of Pyridoxine.HCL; and 0.400 g of Glycine in 875.00 ml of polished D-I H<sub>2</sub>O in sequence. Bring up to volume with polished D-I H<sub>2</sub>O. Make in 400 ml portions. Thiamine.HCL & Pyridoxine.HCL are in Dark Desiccator. Store for one month, unless contamination or precipitation occurs, then make fresh stock.

- 15 Total Volume (L) = 1.00

**560 R**

Ingredient	Amount	Unit
D-I Water, Filtered	950.000	MI
CHU (N6) Basal Salts (SIGMA C-1416)	4.000	G
Eriksson's Vitamin Mix (1000X SIGMA-1511)	4.000	MI
Thiamine.HCL 0.4mg/ml	1.250	MI
Sucrose	30.000	G
2, 4-D 0.5mg/ml	4.000	MI
Gelrite @	3.000	G
Silver Nitrate 2mg/ml #	0.425	MI
Bialaphos 1mg/ml #	3.000	MI

**Directions:**

@ = Add after bringing up to volume

5 # = Add after sterilizing and cooling to temp.

Dissolve ingredients in D-I H<sub>2</sub>O in sequence

Adjust to pH 5.8 with KOH

Bring up to volume with D-I H<sub>2</sub>O

Sterilize and cool to room temp.

10 Total Volume (L) = 1.00

560 Y

Ingredient	Amount	Unit
D-I Water, Filtered	950.000	MI
CHU (N6) Basal Salts (SIGMA C-1416)	4.000	G
Eriksson's Vitamin Mix (1000X SIGMA-1511)	1.000	MI
Thiamine.HCL 0.4mg/ml	1.250	MI
Sucrose	120.000	G
2,4-D 0.5mg/ml	2.000	MI
L-Proline	2.880	G
Gelrite @	2.000	G
Silver Nitrate 2mg/ml #	4.250	MI

## Directions:

@ = Add after bringing up to volume

# = Add after sterilizing and cooling to temp.

- 5 Dissolve ingredients in D-I H<sub>2</sub>O in sequence

Adjust to pH 5.8 with KOH

Bring up to volume with D-I H<sub>2</sub>O

Sterilize and cool to room temp.

\*\* Autoclave less time because of increased sucrose\*\*

- 10 Total Volume (L) = 1.00

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and

15 individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

Applicant's or agent's file reference	5718-72-1	International application No.	PCT/US99/
--	-----------	-------------------------------	-----------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

<b>A.</b> The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 4, line 27	
<b>B. IDENTIFICATION OF DEPOSIT</b> Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depository institution American Type Culture Collection	
Address of depository institution (including postal code and country) 10801 University Blvd. Manassas, Virginia 20110-2209 US	
Date of deposit 21 October 1998	Accession Number 203367
<b>C. ADDITIONAL INDICATIONS</b> (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
Page 4, line 28; Date of deposit 21 October 1998 - Accession Number 203368 Page 4, line 29, Date of deposit 21 October 1998 - Accession Number 203366 Page 4, line 30, Date of deposit 21 October 1998 - Accession Number 203369	
<b>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE</b> (if the indicators are not for all designated States)	
<b>E. SEPARATE FURNISHING OF INDICATIONS</b> (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For International Bureau use only
<input type="checkbox"/> This sheet was received with the International Bureau on:
Authorized officer

## WHAT IS CLAIMED IS:

1. A promoter that drives expression, said promoter comprising a nucleotide sequence selected from the group consisting of:
- a) a nucleotide sequence that drives expression of a homologous nucleotide sequence selected from the group consisting of nucleotide sequences encoding actin-2, enolase, Gos-2 and L41;
  - b) a nucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4;
  - c) a nucleotide sequence comprising a variant or fragment of the nucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4;
  - d) a nucleotide sequence comprising at least 20 contiguous nucleotides of the sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4; and
  - e) a nucleotide sequence that hybridizes to any one of a), b), c), or d) under stringent conditions.
2. The promoter of claim 1, wherein said expression is constitutive.
3. An expression cassette comprising a promoter and a nucleotide sequence operably linked to said promoter, wherein said promoter drives expression of said nucleotide sequence in a plant cell, said promoter comprising a nucleotide sequence selected from the group consisting of:
- a) a nucleotide sequence that drives expression of a homologous nucleotide sequence selected from the group consisting of nucleotide sequences encoding actin-2, enolase, Gos-2 and L41;
  - b) a nucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4;
  - c) a nucleotide sequence comprising a variant or fragment of the nucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4;

d) a nucleotide sequence comprising at least 20 contiguous nucleotides of the sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4; and

e) a nucleotide sequence that hybridizes to any one of a), b), c), or d) under stringent conditions.

4. The expression cassette of Claim 3, wherein expression of said nucleotide sequence operably linked to said promoter alters phenotype of a plant transformed with said expression cassette.

5. A transformation vector comprising the expression cassette of Claim 3.

6. A transformed plant cell having stably incorporated into its genome an expression cassette comprising a promoter and a nucleotide sequence operably linked to said promoter, wherein said promoter drives expression of said nucleotide sequence in a plant cell, said promoter comprising a nucleotide sequence selected from the group consisting of:

a) a nucleotide sequence that drives expression of a homologous nucleotide sequence selected from the group consisting of nucleotide sequences encoding actin-2, enolase, Gos-2 and L41;

b) a nucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4;

c) a nucleotide sequence comprising a variant or fragment of the nucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4;

d) a nucleotide sequence comprising at least 20 contiguous nucleotides of the sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4; and

e) a nucleotide sequence that hybridizes to any one of a), b), c), or d) under stringent conditions.

7. A transformed plant having stably incorporated into its genome an expression cassette comprising a promoter and a nucleotide sequence operably

linked to said promoter, wherein said promoter drives expression of said nucleotide sequence in a plant cell, said promoter comprising a nucleotide sequence selected from the group consisting of:

- 5 a) a nucleotide sequence that drives expression of a homologous nucleotide sequence selected from the group consisting of nucleotide sequences encoding actin-2, enolase, Gos-2 and L41;
- b) a nucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4;
- 10 c) a nucleotide sequence comprising a variant or fragment of the nucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4;
- d) a nucleotide sequence comprising at least 20 contiguous nucleotides of the sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4; and
- 15 e) a nucleotide sequence that hybridizes to any one of a), b), c), or d) under stringent conditions.

8. The plant of claim 7, wherein said plant is a monocot.

20 9. The plant of claim 7, wherein said monocot is maize.

10. The plant of claim 7, wherein said plant is a dicot.

25 11. Seed of the plant of claim 7.


12. The plant of claim 7, wherein said promoter is a maize promoter.

30 13. A method for expressing a nucleotide sequence in a plant cell, said method comprising transforming a plant cell with a transformation vector comprising an expression cassette, said expression cassette comprising a promoter and a nucleotide sequence operably linked to said promoter, wherein said promoter drives expression of said nucleotide sequence in a plant cell, said promoter comprising a nucleotide sequence selected from the group consisting of:



- a) a nucleotide sequence that drives expression of a homologous nucleotide sequence selected from the group consisting of nucleotide sequences encoding actin-2, enolase, Gos-2 and L41;
- b) a nucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4;
- c) a nucleotide sequence comprising a variant or fragment of the nucleotide sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4;
- d) a nucleotide sequence comprising at least 20 contiguous nucleotides of the sequence set forth in SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4; and
- e) a nucleotide sequence that hybridizes to any one of a), b), c), or d) under stringent conditions.
14. A method for expressing a nucleotide sequence in a plant, said method comprising transforming a plant cell according to the method set forth in claim 13, and further comprising regenerating a transformed plant from said plant cell, wherein said plant has stably incorporated into its genome said expression cassette.

Figure 1






Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

## SEQUENCE LISTING

<110> Barbour, Eric  
EuClaire Meyer, Terry  
Eid Saad, Mohammed

<120> Novel Maize Promoters

<130> 5718-72

<160> 4

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 613

<212> DNA

<213> Zea mays

<220>

<221> promoter

<222> (0)...(0)

<223> Actin-2

<400> 1

tcagtatgat	tgctttcaat	ataggctgat	ggagcctatg	aatgatctat	aactatgtga	60
ttggatgtct	taacttgctg	agccaagctc	gtatgagcct	cttttactgg	gtaccactaa	120
tttctatata	ttagatacag	ttaaataagt	ctaaattaca	tcggtgcgtg	cgtgccagta	180
cgactagatg	tagtgaacta	aacacaggtt	aagctgtgat	caatgtatgt	agagagtctg	240
actctttata	tgcggaacaac	ttaacacacg	ataagtgtcg	agaatgattg	aggaaatatt	300
ggcggttcca	cgtgattgag	taagaatgag	agggaataaa	aaggatttgg	tgcgttgggt	360
ttgaaggcag	gatttgggta	ggatgggtgg	acgtttgaag	tgatgagttt	ttcaagcgta	420
tagattttct	atttgcctt	tttaattaac	tttctccag	ccgggatgcg	cgtataaaaa	480
ccggcgaaac	ccttggtct	cctcattcgg	cctatcacaa	ccgcttactc	tcgtgcgctc	540
tccgtgggag	cgaggaccg	cggccggcgg	cagcggcagc	ttctcctaga	tctccggcat	600
catcagtggg	tcc					613

<210> 2

<211> 664

<212> DNA

<213> Zea mays

<220>

<221> promoter

<222> (0)...(0)

<223> Enolase

<400> 2

tcatatattt	atgaacggag	atagtgcac	tatcatgcga	ttcttcgacg	gagtcgacgt	60
ctgattggac	cagaatttca	gatggtgagc	aaaagtgcc	cttgccctgcc	tccttttctc	120
gtgcctgcct	aacacgtgcc	tttgatctct	agggcagtta	taattaagac	aagcaggtta	180
tattataagt	caaggataga	aagaaaggag	aaacaatata	ctatttaaga	tcagataaaa	240
aaagagctaa	taattttttg	ggacacatat	actggttaca	ttgttataat	ctgtatatat	300
cacgttggtc	gaatatattc	caaattttta	ctatgattcg	tgctctaccg	gaactacttc	360
tagattttga	aaactttatg	agaattttct	tatttagata	cactaaggct	aattttgggt	420



ggttttttggc	tcgctagcta	ccattacctc	ctgcatctag	acattacaaa	tttacaataa	480
ataaagttcc	tagattttga	acgaaaccag	cagagcgcac	accgtccttg	ccccacggaa	540
caagaataat	ggaatatgct	cccgcagccc	tcgtggaaac	caagggcgga	ccttcccctc	600
ctccaagcaa	atccgaggcc	cacccacggg	ccgctcgaac	atctgtcacc	gccaaagtgg	660
atcc						664

<210> 3  
 <211> 862  
 <212> DNA  
 <213> Zea mays

<220>  
 <221> promoter  
 <222> (0)...(0)  
 <223> Gos-2

<400> 3						
attattggct	gtaggattct	aaacagagcc	taaatagctg	gaatagctct	atagccctca	60
atccaaacta	atgatattcta	tacttatgca	actctaaatt	tttattctaa	aagtaatat	120
tcatttttgt	caacgagatt	ctctactcta	ttccacaatc	ttttgaagct	atatttacct	180
taaatctgta	ctctatacca	ataatcatat	attctattat	ttatttttat	ctctctccta	240
aggagcatcc	ccctatgtct	gcatggcccc	cgctcgggt	cccaatctct	tgctctgcta	300
gtagcacaga	agaaaacact	agaaatgact	tgcttgactt	agagtatcag	ataaacatca	360
tgtttactta	actttaattt	gtatcggttt	ctactatttt	tataatattt	ttgtctctat	420
agatactacg	tgcaacagta	taatcaacct	tataatattt	ttgtctctat	agatactacg	480
tgcaacagta	taatcaacct	acagcggtcc	ttttatggtt	cctcactggg	cacagcataa	540
acgaaccctg	tccaatgttt	tcggcgcgaa	caaacagaaa	ttccatcagc	gaacaaacaa	600
catacatgcg	agatgaaaat	aaataataaa	aaaagctccg	tctcgatagg	ccggcacgaa	660
tcgagagcct	ccatagccag	ttttttccat	cggaacggcg	gttcgcgcac	ctaattatat	720
gcaccacacg	cctataaagc	caaccaaccc	gtcggagggg	cgcaagccag	acagaagaca	780
gcccgtcagc	ccctctcggt	tttcatccgc	cttcgcctcc	aaccgcgtgc	gctccacgcc	840
tcctccagga	aagcgaggat	cc				862

<210> 4  
 <211> 461  
 <212> DNA  
 <213> Zea mays

<220>  
 <221> promoter  
 <222> (0)...(0)  
 <223> L41

<400> 4						
tctgaaccta	taaaatttgt	tcacatttgt	gtttatcaat	ccactatata	aaagaggatg	60
tagggataa	ctcatgctca	aacctttatt	aaacacatgt	gctagtatct	aggctgctcg	120
atgatgttct	attgttcgcg	actgtgttgt	gtgaaaactg	cagcagatca	actgggccat	180
gctggcctgg	ttatgcacgg	gcacgctcga	agcggaacct	agtgagcagc	ccaacagcgt	240
cctagcggtg	ggcgtacggc	ccattagaag	tacacggccc	gaatccccgg	agaaaaacct	300
agacagcggc	gggcgccttg	ccataaaaagc	acagcgcttc	tacacccgcc	gccgccactc	360
gttgcagcct	ctctacctct	acctaactgga	gtagggagca	cgccgcacga	gccctttccg	420
ccgccgcaa	aatggtgaac	gttcctaaga	ccaagggatc	c		461

